



Jonathan Green

GREEN-UP

Lawn Food with Crabgrass Preventer

PREVENTS CRABGRASS & LISTED ANNUAL WEEDS & GRASSES

FEEDS YOUR LAWN OVER 3 MONTHS

SEASON-LONG CRABGRASS CONTROL

CONTROL CRABGRASS BEFORE IT EMERGES...



AND UP TO 4 WEEKS AFTER IT EMERGES!

20-0-3

ACTIVE INGREDIENT:
Dithiopyr: 3,5-pyridinodicarboximinoic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-5,5-dimethyl ester 0.125%
OTHER INGREDIENTS: 99.875%
TOTAL: 100.000%

This product contains 0.0625 pound of the active ingredient dithiopyr per 50 pound bag.

EPA REG. NO. 62719-483-50931
EPA EST. NO. 075527-PA-12
90561-NC-001 89135-AZ-001
92091-CAN-1 87245-CAN-001
093243-CT-1

Code on bag indicates manufacturing location.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
Refer to back of bag for Directions for Use

USE DIRECTIONS FOR TURF

This turf fertilizer product, containing Dimension® herbicide, can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months of crabgrass control. This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only, but is effective up to 4 weeks after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after crabgrass germination. Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

COOL-SEASON GRASSES		WARM-SEASON GRASSES	
bentgrass, creeping	fescue, fine	bahiagrass	buffalograss
bluegrass, Kentucky	fescue, tall	centipede	carpetgrass
		zoysiagrass	St. Augustinegrass

USE PRECAUTIONS FOR TURF

This product should be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, initial application of this product should be made only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. **Exception:** In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.

- Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrass injury.
- For best results, cultural practices that disturb the soil, such as core, spike, or hydro-aeration, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.
- Reseeding, Overseeding, or Sprigging:** Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 9.2 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

Use Restrictions

- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.

APPLICATION TIMING

SPRING APPLICATION

Control of Crabgrass: This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.

Control or Suppression of Other Grass and Broadleaf Weeds: For control or suppression of other grass and broadleaf weeds, this product must be applied as a preventative treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

FALL APPLICATION

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, do not reseed, overseed or sprig turfgrasses within 12 weeks after application.



TIPS FOR IMPROVED WEED CONTROL

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

APPLICATION INSTRUCTIONS AND EQUIPMENT

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.



BROADCAST (ROTARY) SPREADERS



	2.4 lb. 1,000 sq. ft.	3.2 lb. 1,000 sq. ft.	4.0 lb. 1,000 sq. ft.
Jonathan Green New American Lawn New American Lawn Deluxe & Pro New American Lawn Hand Held	6-1/2 7 14 2	7 13 3 2	7 13 3 2
Agri-Fab Cyclone EZ Republic (Ortho) Earthway (Ames) Precision (Red Devil) Scotts Rotary Scotts EasyGreen	5 4-1/2 3-1/2 15 5 7 26	6 5 4 16 6-1/2 8 28	6 5 4 16 6-1/2 8 28

CALIBRATION SETTINGS AT SPECIFIED RATE

	2.4 lb. 1,000 sq. ft.	3.2 lb. 1,000 sq. ft.	4.0 lb. 1,000 sq. ft.
Drop Spreaders	1-1/2 2 4 14 2 3-1/2	2 7 4-1/2 15 3 4	2 7 4-1/2 15 3 4

DROP SPREADERS



	2.4 lb. 1,000 sq. ft.	3.2 lb. 1,000 sq. ft.	4.0 lb. 1,000 sq. ft.
Agri-Fab Central EZ Republic (Ortho) Earthway (Ames) Precision (Red Devil) Scotts Drop	1-1/2 2 4 14 2 3-1/2	2 7 4-1/2 15 3 4	2 7 4-1/2 15 3 4

GUARANTEED ANALYSIS

Total Nitrogen (N) 20% Derived From: Polymer Coated Urea,
20.00% Water Soluble Nitrogen * Urea, Muriate of Potash.
Soluble Potash (K₂O) 3% *5.0% controlled release urea nitrogen from polymer coated urea

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfa.org/metals.html>

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes Eye Irritation • Harmful If Inhaled
Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

- IF IN EYES** • Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.
- IF INHALED** • Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Notifies Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.
In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:
1. Refund of purchase price paid by buyer or user for product bought; or
2. Replacement of amount of product used.
To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, do not exceed 4.6 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre). In the state of New York, this product may be applied at a minimum of 2.4 lb per 1000 sq ft per year (equivalent to 0.128 lb active ingredient per acre).



®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

