

Jonathan Green®

# VERI-GREEN®


## Lawn Food with Crabgrass Preventer

PREVENTS  
CRABGRASS & LISTED  
WEEDS & GRASSES

FEEDS YOUR  
LAWN FOR UP TO  
3 MONTHS

SEASON LONG  
CRABGRASS  
CONTROL

CONTROL CRABGRASS  
BEFORE  
IT EMERGES...



AND UP TO 4 WEEKS  
AFTER  
IT EMERGES!

20-0-3

**ACTIVE INGREDIENT:**  
Dithiopyr: 3,5-pyridinedicarboxylic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-  
(trifluoromethyl)-5,5-dimethyl ester ..... 0.125%  
**OTHER INGREDIENTS:** ..... 99.875%  
TOTAL: ..... 100.000%

EPA REG. NO. 62719-483-50931  
EPA EST. NO. 075527-PA-12  
90561-NC-001 89135-AZ-001  
093243-CI-1 8378-IN-1  
69759-OR-2 69759-OR-004 (LOT BEGINS BAN)

KEEP OUT OF REACH  
OF CHILDREN  
**CAUTION**  
Refer to back of bag for  
Directions for Use

- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, do not exceed 4.6 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre). In the state of New York, this product may be applied at a minimum of 2.4 lb per 1000 sq ft per year (equivalent to 0.128 lb active ingredient per acre).

### GUARANTEED ANALYSIS

Total Nitrogen (N) .....	20%
20.00% Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> O) .....	3%
Derived From: Polymer Coated Urea, Urea, Muriate of Potash.	
*5.0% controlled release urea nitrogen from polymer coated urea	
Information regarding the contents and levels of metals in this product is available on the internet at <a href="http://www.aapfo.org/metals.html">http://www.aapfo.org/metals.html</a>	

### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

**Causes Eye Irritation • Harmful if Inhaled**

**Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.**

### ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**Pesticide Storage:** Store this product only in its original container in a dry, cool, secured storage area.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

**If empty:** Place in trash or offer recycling if available.

**If partially filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

### USE DIRECTIONS FOR TURF

This turf fertilizer product, containing Dimension® herbicide, can be applied 4 weeks after preemergent crabgrass timing and provides up to 4 months of crabgrass control.

This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only, but is effective up to 4 weeks after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after crabgrass germination.

**Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:**

COOL-SEASON GRASSES:			WARM-SEASON GRASSES:		
Bentgrass, creeping	Fescue, fine	Ryegrass, perennial	Bahia grass	Buffalograss	Centipedegrass
Bluegrass, Kentucky	Fescue, tall		Bermudagrass	Carpetgrass	St. Augustinegrass
					Zoysiagrass

### USE PRECAUTIONS FOR TURF

This product should be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, initial application of this product should be made only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. **Exception:** In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.

- Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrass injury.
- For best results, cultural practices that disturb the soil, such as core, spike, or hydro-aerification, and verticutting, should be done before application of this product. **Do not** apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.
- **Reseeding, Overseeding, or Sprigging:** Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 9.2 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

### USE RESTRICTIONS

- **Do not** use clippings from treated turf for mulching around vegetables or fruit trees.
- **Do not** allow people or pets to enter treated area until dust has settled.

### APPLICATION TIMING

#### SPRING APPLICATION

**Control of Crabgrass:** This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.

**Control or Suppression of Other Grass and Broadleaf Weeds:** For control or suppression of other grass and broadleaf weeds, this product must be applied as a **preventative** treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

#### FALL APPLICATION

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, **do not** reseed, overseed or sprig turfgrasses within 12 weeks after application.

#### TIPS FOR IMPROVED WEED CONTROL

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

### WEEDS CONTROLLED OR SUPPRESSED

This product will provide control of crabgrass and control or suppression of other listed grass and broadleaf weeds when applied prior their germination.

GRASSES			BROADLEAF WEEDS		
barley	goosegrass	bittercress ↑	medic, black	shepherdspurse	
barnyardgrass	foxtail, green	carpetweed	mustard	speedwell, corn ↑	
bluegrass, annual	foxtail, yellow	chickweed ↑	oxalis, buttercup	spurge, garden	
brome	kikuyugrass ↑	geranium, Carolina ↑	pineappleweed	spurge, prostrate	
crabgrass, large	oats, wild	henbit	pigweed, redroot	spotted	
crabgrass, smooth	ryegrass, (annual &	knotweed, prostrate	parsley-piert ↑	woodsorrel, creeping	
crabgrass, Southern	perennial	lespedeza, common	purslane, common	woodsorrel, yellow	
crowfootgrass ↑	sandbur	marestail			
dallisgrass	smutgrass				

↑ Suppression only

### LENGTH OF CONTROL

This product will provide up to 4 months of crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. Refer to the map below to determine the region of the U.S. that includes your location. Use the table below to determine the amount of this product to apply in your location based on the desired length of weed control or need for postemergence crabgrass control.

Use "Regions" within the United States:

☐ North  
☐ South



Amount of this Product to Apply (Pounds of product to apply per 1,000 square feet of turfgrass†)			
--- Length of Crabgrass Control ---			
Region of U.S.	2-3 Months	3-4 Months	Early Postemergence Control
North	2.4 lb/1000 sq ft (0.128 lb ai/acre)	3.2 lb/1000 sq ft (0.174 lb ai/acre)	4.0 - 9.2 lb/1000 sq ft †† (0.22 - 0.5 lb ai/acre)
South	3.5 lb/1000 sq ft (0.192 lb ai/acre)	4.0 - 4.7 lb/1000 sq ft †† (0.22-0.256 lb ai/acre)	4.0 - 9.2 lb/1000 sq ft †† (0.22 - 0.5 lb ai/acre)

† 1,000 square feet equals an area 100 feet X 10 feet or square area of about 32 feet on a side.  
†† Use the higher rate in rate range in areas of high weed pressure or for increased length of control.

#### Maximum Application Rates:

- **Do not** apply more than 6 times per year and do not apply more than 9.2 lb of this product per 1000 sq ft (0.5 lb ai/acre) per application.
- **Do not** apply more than 27.6 lb of this product/1000 sq ft per year (equivalent to 1.5 lb of active ingredient per acre, respectively).
- In the state of New York, do not exceed 4.6 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre).
- To minimize the potential for plant injury, do not make additional applications within five weeks of previous application.

### APPLICATION INSTRUCTIONS AND EQUIPMENT

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides.

For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the specified application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area.

More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.

\*These settings are approximate for the application rate shown per 1,000 square feet. You may need to adjust the spreader setting depending on walking speed, spreader condition and product.

### CALIBRATION SETTINGS AT SPECIFIED RATE

BROADCAST (ROTARY) SPREADERS		3.2 lb, 1,000 sq. ft.	DROP SPREADERS		3.2 lb, 1,000 sq. ft.
Jonathan Green Standard Deluxe & Pro Hand Held		5-1/2 14 4 5	Central		6
Agri-Fab		4	EZ Republic (Ortho)		4
Chapin 81000A		5	Earthway (Ames)		14
Cyclone		4-1/2	Precision (Red Devil)		5
EZ Republic (Ortho)		9	Scotts Drop**		7
Earthway (Ames)		14			
Lesco		11 - 12 / D - E			
Precision (Red Devil)		5			
Scotts Rotary*		5			
Scotts EasyGreen		26			

\*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 3000

### TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

### WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

### INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

### LIMITATION OF REMEDIES

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Dimension®  
SPECIALTY HERBICIDE

®Trademark of The Dow  
Chemical Company  
("Dow") or an affiliated  
company of Dow

MADE IN USA




26000 59122







GREAT  
Lawns ONLY  
Happen with  
  
PROPER  
Soil pH...  
  
and  
LOOSE,  
Biologically  
Active Soil!



**Jonathan Green**  
QUALITY LAWNS SINCE 1881

# MAG-I-CAL PLUS

**Soil Food**  
for Lawns in Alkaline + Hard Soils

HELPS  
LOWER  
SOIL pH

LOOSENS  
HARD  
SOILS

INCREASES  
POROSITY OF  
SOILS




LAWN FOOD NUTRIENTS  
ARE WASTED WHEN THE  
SOIL pH IS NOT BALANCED!

PROMOTES AIR AND  
WATER PENETRATION  
FOR BETTER ROOTING

## THE NEW AMERICAN LAWN

FEED YOUR LAWN  
...AND YOUR SOIL!



### EARLY SPRING:

Apply Green-Up  
plus Crabgrass  
Preventer

### LATE SPRING:

Apply Green-Up  
Weed & Feed

### SUMMER:

Apply  
Mag-I-Cal Plus

### FALL:

Apply Winter  
Survival

\*For very hard, compact soil, supplement this program with Love Your Soil. See right for more details.

### FEEDING YOUR SOIL

If you've struggled to grow a great lawn in the past, poor soil is often the problem. If your soil is too alkaline (having a pH over 7) or compacted, your lawn may not thrive. MAG-I-CAL PLUS for Alkaline Soils contains calcium which improves cell wall structure. Lawns thrive in soil with pH levels between 6.2 and 7; otherwise light green grass and weeds may persist, while fertilizer nutrients are not fully used by the grass. MAG-I-CAL PLUS helps to break up clay and compacted soils, which improves root penetration and promotes growth by increasing biological activity while helping to release trapped nutrients in the soil.

### WHY USE SOIL FOOD?

**HARD SOIL\***: For hard, compact soil, supplement our New American Lawn Program with LOVE YOUR SOIL.

Apply any time of the year with other products on the same day. It can be applied when seeding or not.

### DIRECTIONS FOR USE

MAG-I-CAL PLUS for alkaline soils can be applied anytime of the year. Apply only as directed.

**ESTABLISHED LAWNS:** Apply MAG-I-CAL PLUS using the "Spreader Settings" chart below. For best results, aerate or rake the lawn vigorously prior to applying MAG-I-CAL PLUS. Watering is not necessary, but if possible, water in after application for 15 minutes. If soil is above 7.5, apply again in 3 months.

**NEW LAWNS OR OVERSEEDING LAWNS OR SODDING:** When seeding or sodding lawns, MAG-I-CAL PLUS should be applied with Jonathan Green Green-Up Lawn Food for Seeding & Sodding. These products can be applied in any order, on the same day.

Apply MAG-I-CAL PLUS using the "Spreader Settings" chart below. For best results, aerate or rake the lawn vigorously prior to applying MAG-I-CAL PLUS, grass seed or any of the Jonathan Green products mentioned above. Watering is not necessary, but if possible, water in after application for 15 minutes.




### SPREADER SETTINGS

The following settings are approximate. An accurate calculation can be made by application to 1,000 sq. ft. and adjusting spreader opening to apply 3.6 lbs. per 1,000 sq. ft. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .90 pounds per 250 square feet. Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

#### BROADCAST (ROTARY) SPREADERS

	Established lawns	New lawns or pH over 7.5 covers
	4-1/2	Apply
	13	twice over
	3	the area
Agri-Fab	2-1/2	at the
Cyclone	4	same
EZ Republic (Ortho)	4	setting to
Earthway (Ames)	12	the left
Precision (Red Devil)	5	
Scotts Rotary	5	
Scotts EasyGreen	20	

#### DROP SPREADERS

	Established lawns 5,000 sq. ft.	New lawns or pH over 7.5 covers 2,500 sq. ft.
	2	Apply
Central	4	twice over
EZ Republic (Ortho)	4	the area
Earthway (Ames)	10	at the
Precision (Red Devil)	3-1/2	same
Scotts	1-1/2	setting to the left

For more settings visit [www.jonathangreen.com](http://www.jonathangreen.com)

### USAGE TIPS

#### For rectangular lawns


Apply two header strips across each end for a turning area. Then apply back and forth in the longest direction.

#### For irregular-shaped lawns

Apply a header strip around the entire lawn for a turning area, then apply back and forth.

**Note:** For Rotary Spreaders each pass should be within 10 feet of each other to ensure an overlap.

- Do not leave spreader open when making turns.
- For best results, do not apply on a windy day.
- Swath will vary due to weather conditions, product used and walking speed.
- Apply products only to lawn.
- To determine how much coverage you need, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, gardens, walkways, etc.).



### GUARANTEED ANALYSIS

Calcium (Ca)	18.0%
Sulfur (S)	17.0%
17% Combined Sulfur (S)	

Derived from: Calcium Sulfate Dihydrate, Ammonium Sulfate and Sulfuric Acid.

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:  
6% Humic Acid (derived from humus)  
1% Polyhydroxycarboxylic Acids (derived from plant extracts)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

### CAUTION

Store in a cool dry place out of the reach of children and pets. Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn or garden and sweep any product that lands on the driveway, sidewalk or street back onto your lawn or garden. Also, sweep fertilizer off concrete surfaces to avoid staining.

### NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all risk of use and/or handling if such use is contrary to label instructions.

### CAS NO.

Calcium Sulfate Dihydrate	10101-41-A
Humic Acid	1415-93-6
Sulfuric Acid	7664-93-9
Ammonium Sulfate	7783-20-2

Be aware of local ordinance requirements concerning lawn fertilizer applications. "Black out" dates may limit or prohibit lawn fertilizer applications from late fall to early spring in certain areas.

### My Pledge Of Satisfaction

We have made every effort in the formulation of this product to insure that you will be satisfied with the results. If, for any reason, you are not satisfied, please write and describe the problem. Our lawn specialists will make every effort to assure your success.



Jonathan Green



Sign up to receive timely lawn tips  
[www.jonathangreen.com](http://www.jonathangreen.com)

Distributed & Guaranteed by: Jonathan Green, Inc. POB 326 Farmingdale, NJ 07727





**Jonathan Green**  
QUALITY LAWN SINCE 1881

# WINTER SURVIVAL

## Fall Lawn Food

HELPS REPAIR  
LAWN FROM  
SUMMER DAMAGE

IDEAL FOR  
FALL  
SEEDING

HELPS PROTECT  
LAWN FROM  
WINTER STRESS

PROMOTES DEEP ROOTING IN FALL...  
AND A QUICKER GREEN-UP IN SPRING!

**10-0-20**

### FALL FEEDING

Fall is a great time to grow grass and feed your lawn. Cooler temperatures and higher amounts of rainfall helps your lawn recover from summer stress. Winter Survival Fall Lawn Food provides nutrients to make your lawn thicker and greener than ever before while encouraging deeper-rooting. Winter Survival Fall Lawn Food can be used on all grass types.

### HOW TO APPLY

Apply Winter Survival Fall Lawn Food from late summer through fall. For damaged lawns, apply in early fall and again 6 weeks later for a thicker, greener lawn. Apply when grass is dry and water thoroughly after applying for best results; however, product does not require watering in. You may apply MAG-I-CAL Plus in any order on the same day that you apply Winter Survival Fall Lawn Food. You can re-enter lawn area after product is applied. Apply uniformly with a lawn spreader using the chart listed. This package applies 3 lb. of product per 1,000 square feet.

### SEEDING LAWNS

Winter Survival Fall Lawn Food is ideal for fall seeding. Apply Winter Survival Fall Lawn Food after grass seed has been planted. You may apply MAG-I-CAL Plus in any order on the same day that you apply Winter Survival Fall Lawn Food. Water the lawn after application to activate the feeding process and get new grass plant seedlings off to a good start.

### CARING FOR YOUR LAWN

Mow grass at 2-1/2 to 3-1/2 inches. Do not remove more than 1/3 of the grass blade at one time, with a sharp mower blade. Mulch clippings or leave them on the lawn if not wet or clumpy. One inch of water from rainfall and/or watering per week is usually adequate unless drought conditions exist.

### SPREADER SETTINGS

The following settings are approximate for applying contents of this package at a suggested rate of 3 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .75 pounds per 250 square feet. Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

#### BROADCAST (ROTARY) SPREADERS

<b>Jonathan Green</b>	<b>5</b>
<b>Standard</b>	<b>10-11</b>
<b>Deluxe &amp; Pro</b>	<b>3</b>
<b>Hand Held</b>	
Agri-Fab	6
Cyclone	5
Earthway (Ames)	14
Lesco	12 - E
Ortho	5
Precision (Red Devil)	5
Scotts Rotary Spreaders	6*
Scotts EasyGreen	26


\*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 3000

#### DROP SPREADERS

Earthway (Ames)	14-1/2
Ortho	5
Precision (Red Devil)	3
Scotts Drop**	7


\*\*Includes Accugreen 1000 - 2000 & Classic

### USAGE TIPS



#### For rectangular lawns

Apply two header strips across each end for a turning area. Then apply back and forth in the longest direction.



#### For irregular-shaped lawns

Apply a header strip around the entire lawn for a turning area, then apply back and forth.

**Note:** Each pass should be within 10 feet of each other to ensure an overlap.

- Do not leave spreader open when making turns.
- For best results, do not apply on a windy day.
- Swath will vary due to weather conditions, product used and walking speed.
- To determine how much coverage you need, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, gardens, walkways, etc.).

### GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	10.00%
10% Urea Nitrogen*	
SOLUBLE POTASH (K <sub>2</sub> O)	20.00%
IRON (Fe)	3.00%
0.03% water soluble iron	

Derived from: Urea, Polymer Coated Urea, Muriate of Potash, Ferrous Sulfate.

\*2.5% Slowly available urea nitrogen from polymer coated urea.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

### CAUTION

Keep out of reach of children. Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn or garden and sweep any product that lands on the driveway, sidewalk or street back onto your lawn or garden. Sweep fertilizer off concrete surfaces to avoid staining.

### NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling if such use is contrary to label instructions.

**Be aware of local ordinance requirements concerning lawn fertilizer applications. "Black out" dates may limit or prohibit lawn fertilizer applications from late fall to early spring in certain areas.**

#### My Pledge Of Satisfaction

We have made every effort in the formulation of this product to assure that you will be satisfied with the results. If, for any reason, you are not satisfied, please write and describe the problem. Our lawn specialists will make every effort to assure your success.

Jonathan Green

Ask for a free copy of our  
Lawn Guide.

Sign up to receive timely lawn tips  
[www.jonathangreen.com](http://www.jonathangreen.com)



Distributed & Guaranteed by: Jonathan Green, Inc. POB 326 Farmingdale, NJ 07727

(22400) SF621