

Jonathan Green


LAWN WEED CONTROL

Broadleaf Weed Control

APPLY
SPRING - SUMMER - FALL

CONTROLS DANDELION
AND OVER 200 LISTED
BROADLEAF WEEDS

WILL NOT
BURN YOUR
LAWN



DANDELION

WHITE CLOVER

CHICKWEED

OXALIS

ACTIVE INGREDIENTS	
2,4-dichlorophenoxyacetic acid	1.37%
(+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.31%
Dicamba: 3,6-dichloro-o-anisic acid	0.13%
OTHER INGREDIENTS	
TOTAL	98.19%
	100.00%

THIS PRODUCT CONTAINS:
27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
6.2 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
2.6 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
Isomer Specific by ADAC Methods
TRIMEC® is a registered trademark of PBI/Gordon Corporation

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for complete Precautionary Statements and Directions for Use, including First Aid and Storage and Disposal.
EPA REG. NO. 2217-640-50931 EPA EST. NO. 075527-PA-12
90561-NC-001 89135-AZ-001 92091-CAN-1
87245-CAN-001 093243-CT-1 69759-OR-004
Code on bag indicates manufacturing location.

STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Causes moderate eye irritation. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes, and gloves. Remove and wash contaminated clothing before reuse.

FIRST AID

If on skin or on clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- For residential use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Apply when the air is calm. Do not apply to newly seeded lawns. Do not exceed specified dosages for any area. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS: Jonathan Green Lawn Weed Control controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others as listed which are commonly found in home lawns. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days.

For use on established turf of warm season grasses including bermudagrass, centipedegrass, and zoysiagrass. Established turf of cool season grasses including Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra, St. Augustinegrass, or established turf containing desirable clovers.

The maximum broadcast application rate is 2.0 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 4.0 pounds of product per 1,000 sq. ft.

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Controls over 200 weeds!

WEEDS	
Annual yellow sweetclover	Catnip
Aster	Catsear
Austrian fieldcress	Chickweed, common
Bedstraw	Chickweed, mouseear
Beggarticks	Chicory
Betony, Florida	Cinquefoil
Bindweed, field	Clover, crimson
Bird vetch	Clover, hop
Bitter wintercress	Clover, red
Bittercress, hairy	Clover, strawberry
Bitterweed	Clover, sweet
Black-eyed Susan	Clover, white
Black medic	Cockle
Black mustard	Cocklebur
Blackseed plantain	Common mullein
Blessed thistle	Creeping jenny
Bloodflower milkweed	Cudweed
Blue lettuce	Curly dock
Blue vervain	Daisy, English
Bracted plantain	Daisy, fleabane
Brassbuttons	Daisy, oxeye
Bristly oxtongue	Dandelion
Broadleaf dock	Dichondra
Broadleaf plantain	Dogbane
Broomweed	Dogfennel
Buckhorn	Dollarweed
Buckhorn buttercup	False dandelion
Bulbous plantain	False sunflower
Bull thistle	Fiddleneck
Bullnettle	Florida pusley
Burclover	Frenchweed
Burdock	Galinsoga
Burning nettle	Goldenrod
Burweed	Ground ivy
Buttercup	Gumweed
Buttonweed	Hairy fleabane
Canada thistle	Hawkweed
Carolina geranium	Healall
Carpetweed	Heartleaf drymary
Catchweed bedstraw	Heath aster
	Hedge bindweed
	Hedge mustard
	Hemp
	Henbit
	Hoary cress
	Hoary plantain
	Hoary vervain
	Horsenettle
	Jimsonweed
	Knawel
	Knotweed
	Kochia
	Lambsquarters
	Lespedeza
	Mallow
	Matchweed
	Mexicanweed
	Milk vetch
	Morningglory
	Mouseear hawkweed
	Mugwort
	Musk thistle
	Mustard
	Narrowleaf plantain
	Narrowleaf vetch
	Nettle
	Orange hawkweed
	Oriental cocklebur
	Oxalis
	Parsley-piert
	Parsnip
	Spiny amaranth
	Spiny cocklebur
	Spiny sowthistle
	Spotted catsear
	Spotted spurge
	Spurweed
	Stinging nettle
	Strawberry,
	India mock
	Tall nettle
	Tall vervain
	Tansy ragwort
	Tansy mustard
	Tanweed
	Thistle
	Trailing crownvetch
	Tumble mustard
	Tumble pigweed
	Velvetleaf
	Venice mallow
	Virginia buttonweed
	Virginia creeper
	Virginia pepperweed
	Wavyleaf bullthistle
	Western clematis
	Wild buckwheat
	Wild carrot
	Wild four-o'clock
	Wild garlic
	Wild geranium
	Wild lettuce
	Wild marigold
	Wild mustard
	Wild onion
	Wild parsnip
	Wild radish
	Wild rape
	Wild strawberry
	Wild sweet potato
	Wild vetch
	Woodsorrel
	Woolly croton
	Woolly morningglory
	Woolly plantain
	Wormseed
	Yarrow
	Yellow rocket
	Yellowflower pepperweed

SPREADER SETTINGS

The spreader settings below are approximate. Calibrate your spreader on 100 sq. ft. and adjust to apply at the rate of 1 pound of product per 500 sq. ft. (2 lbs. of material per 1,000 sq. ft.). Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.

ROTARY SPREADERS


Jonathan Green	
New American Lawn Standard	4-1/2
New American Lawn Deluxe & Pro	11
New American Lawn Hand Held	3
Agri-Fab	3 - 4
Chapin	15
Cyclone	4
Earthway	
ACE - Greenthumb	
Agway - Fertilome	16
EZ Republic (Ortho)	2
Greenview Jet-Spred SS1	16
LESCO (lettered & numbered)	G - 15
Precision Greenline (Red Devil)	5
Scotts Rotary Spreaders	5 - 6
Scotts EasyGreen	26
Vigaro	3

DROP SPREADERS

Agri-Fab	4
Central 550	8 - 9
Earthway (Ames)	12
EZ Republic (Ortho)	5-1/2
Lawn Crafter	5
Precision Greenline (Red Devil)	4
Scotts Drop Spreaders	4 - 5
Spyker Pro	33



Apply in a single pass at a rate of 2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off the spreader before making turns.



Apply in a single pass at a rate of 2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off the spreader before making turns.

***IMPORTANT:** It is important to follow the directions as indicated on this quality lawn care product as over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

Spreading Hints: These granular settings are approximate. Calibrate your spreader on 100 square feet and adjust to apply at the rate of 1 pound of product per 500 square feet. This bag covers 5,000 sq. ft. of lawn area. To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.

Distributed by: Jonathan Green, Inc. POB 326 Farmingdale, NJ 07727