

Jonathan Green

VERI-GREEN®

Lawn Food with Crabgrass Preventer

PREVENTS CRABGRASS & LISTED WEEDS & GRASSES

FEEDS YOUR LAWN FOR UP TO 3 MONTHS

SEASON LONG CRABGRASS CONTROL

CONTROL CRABGRASS BEFORE IT EMERGES...



AND UP TO 4 WEEKS AFTER IT EMERGES!

20-0-3

ACTIVE INGREDIENT:
Dithiopyr: 3,5-pyridinedicarboxylic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-5,5-dimethyl ester 0.125%
OTHER INGREDIENTS: 99.875%
TOTAL: 100.000%

EPA REG. NO. 62719-483-50931
EPA EST. NO. 075527-PA-12
90561-NC-001 89135-AZ-001
093243-CT-1 8378-IN-1
69759-OR-2 69759-OR-004 (LOT BEGINS BAN)
Code on bag indicates manufacturing location.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
Refer to back of bag for Directions for Use

- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, do not exceed 4.6 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre). In the state of New York, this product may be applied at a minimum of 2.4 lb per 1000 sq ft per year (equivalent to 0.128 lb active ingredient per acre).

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
20.00% Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	3%
Derived From: Polymer Coated Urea, Urea, Muriate of Potash.	
*5.0% controlled release urea nitrogen from polymer coated urea	
Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfo.org/metals.html	

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Causes Eye Irritation - Harmful if Inhaled
Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.
Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened. In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.
Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area.
Container Handling: Nonrefillable container. Do not reuse or refill this container.
If empty: Place in trash or offer recycling if available.
If partially filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE DIRECTIONS FOR TURF

This turf fertilizer product, containing Dimension® herbicide, can be applied 4 weeks after preemergent crabgrass timing and provides up to 4 months of crabgrass control. This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only, but is effective up to 4 weeks after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after crabgrass germination.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

COOL-SEASON GRASSES:			WARM-SEASON GRASSES:		
Bentgrass, creeping	Fescue, fine	Ryegrass, perennial	Bahiagrass	Buffalograss	Centipedegrass
Bluegrass, Kentucky	Fescue, tall		Bermudagrass	Carpetgrass	St. Augustinegrass
					Zoysiagrass

USE PRECAUTIONS FOR TURF

- This product should be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, initial application of this product should be made only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. **Exception:** In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.
- Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrass injury.
 - For best results, cultural practices that disturb the soil, such as core, spike, or hydro-aerification, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.
 - **Reseeding, Overseeding, or Sprigging:** Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 9.2 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

USE RESTRICTIONS

- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.

APPLICATION TIMING

SPRING APPLICATION
Control of Crabgrass: This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.

Control or Suppression of Other Grass and Broadleaf Weeds: For control or suppression of other grass and broadleaf weeds, this product must be applied as a preventative treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

FALL APPLICATION

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, do not reseed, overseed or sprig turfgrasses within 12 weeks after application.

TIPS FOR IMPROVED WEED CONTROL

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

WEEDS CONTROLLED OR SUPPRESSED

This product will provide control of crabgrass and control or suppression of other listed grass and broadleaf weeds when applied prior their germination.

GRASSES		BROADLEAF WEEDS	
barley	goosegrass	bittercrss †	medic, black
barnyardgrass	foxtail, green	carpetweed †	mustard
bluegrass, annual	foxtail, yellow	chickweed †	oxalis, buttercup
brome	kikuyugrass †	geranium, Carolina †	pineappleweed
crabgrass, large	oats, wild	henbit	parsley-piert †
crabgrass, smooth	ryegrass, (annual & perennial)	knotweed, prostrate	purslane, common
crabgrass, Southern	sandbur	lespedeza, common	rocket, London
crowfootgrass †	smutgrass	marestail	
dallisgrass			

† Suppression only

LENGTH OF CONTROL

This product will provide up to 4 months of crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. Refer to the map below to determine the region of the U.S. that includes your location. Use the table below to determine the amount of this product to apply in your location based on the desired length of weed control or need for postemergence crabgrass control.

Use "Regions" within the United States:



Region of U.S.	Amount of this Product to Apply (Pounds of product to apply per 1,000 square feet of turfgrass)†		
	--- Length of Crabgrass Control ---		
North	2-3 Months (0.128 lb ai/acre)	3-4 Months (0.174 lb ai/acre)	Early Postemergence Control (4.0 - 9.2 lb/1000 sq ft †† (0.22 - 0.5 lb ai/acre)
	South	3.5 lb/1000 sq ft (0.192 lb ai/acre)	4.0 - 4.7 lb/1000 sq ft †† (0.22-0.256 lb ai/acre)

† 1,000 square feet equals an area 100 feet X 10 feet or square area of about 32 feet on a side.
†† Use the higher rate in rate range in areas of high weed pressure or for increased length of control.

Maximum Application Rates:

- Do not apply more than 6 times per year and do not apply more than 9.2 lb of this product per 1000 sq ft (0.5 lb ai/acre) per application.
- Do not apply more than 27.6 lb of this product/1000 sq ft per year (equivalent to 1.5 lb of active ingredient per acre, respectively).
- In the state of New York, do not exceed 4.6 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre).
- To minimize the potential for plant injury, do not make additional applications within five weeks of previous application.

APPLICATION INSTRUCTIONS AND EQUIPMENT

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the specified application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area.

More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application. Check equipment frequently to verify calibration and proper functioning.

*These settings are approximate for the application rate shown per 1,000 square feet. You may need to adjust the spreader setting depending on walking speed, spreader condition and product.

CALIBRATION SETTINGS AT SPECIFIED RATE

BROADCAST (ROTARY) SPREADERS	3.2 lb. 1,000 sq. ft.	DROP SPREADERS	3.2 lb. 1,000 sq. ft.
Jonathan Green Standard	5-1/2	Central	6
Deluxe & Pro	14	EZ Republic (Ortho)	4
Hand Held	4	Earthway (Ames)	14
Agri-Fab	5	Precision (Red Devil)	5
Chapin 81000A	5	Scotts Drop**	7
Cydnone	4-1/2		
EZ Republic (Ortho)	9		
Earthway (Ames)	14		
Lesco	11 - 12 / D - E		
Precision (Red Devil)	5		
Scotts Rotary*	5		
Scotts EasyGreen	26		

*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 3000

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Dimension®
SPECIALTY HERBICIDE



®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

MADE IN USA

0 79545 16000 5

VERI-GREEN CRABGRASS

26000 51922



Jonathan Green

VERI-GREEN®

Weed & Feed

**FEEDS LAWNS
FOR UP TO
3 MONTHS**

**KILLS WEEDS
DOWN TO
THE ROOT**

**FOR USE
IN SPRING
OR FALL**




DANDELION



CLOVER

**CONTROLS
250+
BROADLEAF
WEEDS!**



CHICKWEED



OXALIS

21-0-3

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid	0.700%
Dimethylamine Salt of (+)-8-(2-(2-Methyl-4-Chlorophenoxy)propionic Acid)	0.182%
Dimethylamine Salt of (+)-8-(2-(2,4-Dichlorophenoxy)propionic Acid)	0.179%
OTHER INGREDIENTS:	98.939%
TOTAL:	100.000%

DISSOLVE® is a Registered Trademark of Nufarm Americas Inc. Patent #5,266,553

ISOMER SPECIFIC METHOD, EQUIVALENT TO:

**2,4-Dichlorophenoxyacetic Acid	0.58%, 0.0058 oz/lb
**(+)-8-(2-(2-Methyl-4-Chlorophenoxy)propionic Acid)	0.15%, 0.0015 oz/lb
**(-)-8-(2-(2,4-Dichlorophenoxy)propionic Acid)	0.15%, 0.0015 oz/lb

†CONTAINS THE SINGLE ISOMER FORMS OF MECPROPO-P and DICHLOROPROP.
EPA REG. NO. 228-280-50931
EPA EST. NO. 075527-PA-12, 05561-NC-001, 81135-AZ-001, 092343-CF-1, 9198-OH-1M, 9198-OH-2B, 9198-AL-001A, 82757-OH-001, 8378-IN-1, 66759-OR-2, 66759-OR-004 (LOT BEGINS BAN)
Code on bag indicates manufacturing site.

KEEP OUT OF REACH OF CHILDREN

CAUTION
SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS
For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

**THE NEW AMERICAN LAWN
FEED YOUR LAWN
...AND YOUR SOIL!**



EARLY SPRING:
Apply Veri-Green plus
Crabgrass Preventer

LATE SPRING:
Apply Veri-Green
Weed & Feed

SUMMER:
Apply MAG-ICAL Plus

FALL:
Apply Winter Survival

Ask for a FREE copy of our
Lawn Care Guide or go to
www.jonathangreen.com

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND LAWN FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION. WEEDS CONTACTED BY THIS PRODUCT GRADUALLY DIE AND DISAPPEAR FROM THE LAWN.

WHEN TO APPLY VERI-GREEN™ WEED & FEED

Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (pinpoint) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results: (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possibilities retreating in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses.

It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

SPREADER SETTINGS (One pass application)

The following settings are approximate for applying contents of this package at a suggested rate of 3 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .75 pounds per 250 square feet. Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

BROADCAST (ROTARY) SPREADERS

Jonathan Green Standard	3-1/2
Deluxe & Pro Hand Held	11
Agri-Fab	2
Chopin 81000A	4-1/2
Cyclone	4-1/2
EZ Republic (Ortho)	3-1/2
Earthway (Ames)	13
Lesco	10 - 11/D
Precision (Red Devil)	4
Scotts Rotary Spreaders*	4-1/2
Scotts EasyGreen	24

*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 3000

DROP SPREADERS

Central	4
EZ Republic (Ortho)	3
Earthway (Ames)	13
Precision (Red Devil)	3
Scotts**	4

**Includes Accugreen 1000 - 2000 & Classic

For more settings visit
www.jonathangreen.com

GUARANTEED ANALYSIS:

Total Nitrogen (N)	21%
21.00% Urea Nitrogen*	
Soluble Potash (K ₂ O)	3%
Derived From: Polymer Coated Urea, Urea, Sulfate of Potash	
*6.30% slowly available urea nitrogen from polymer coated urea	

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.cnpfa.org/veridb.html>

FIRST AID

- IF ON SKIN OR CLOTHING**
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15 to 20 minutes.
 - Call a poison control center or doctor for treatment advice.
- IF INHALED**
- Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible.
 - Call a poison control center or doctor for further treatment advice.
- IF IN EYES**
- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- HOT LINE NUMBER**
- Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in waters adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run-off to water bodies or drainage systems.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fishponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area to the point of run-off or apply when raining or when rain is expected that day.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material. Wash thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.

USE PRECAUTIONS FOR TURF, ORNAMENTAL (golf courses, cemeteries, parks, sports fields, turfgrass, lawns and other grass areas)

Postemergence, limited to 2 applications per year. Maximum of 250 pounds of product/acre per application. The maximum seasonal rate is 500 pounds of product/acre, excluding spot treatments.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing 63 pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

HOW TO APPLY VERI-GREEN™ WEED & FEED

This 15 pound bag should be used on 5,000 square feet or apply at 3 pounds per 1,000 square feet. Apply evenly with a lawn spreader as a one pass application. **DO NOT HAND SPREAD.**

DIRECTIONS FOR USE



It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS. A minimum of 30 days between applications is required. This product is a stable mixture of three herbicides. It is for use on all residential turf. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dust has settled. This product may be used on Bahia, common Bermudagrass, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine, and Zoysia grass. **DO NOT USE** on Dichondra and Carpet grass. **DO NOT USE** this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

- | | | | |
|-----------------------|-------------------------|-------------------|-------------------------|
| Alder | Clover, White | Hemp | Onion, Wild |
| Antelope | Cockle | Hemlock | Oxalis |
| Aster | Cocklebur | Hoary Cress | Pansy, Field |
| Aster, Wild | Cocklebur, Spiny | Honeysuckle | Parsley-piert |
| Bedstraw | Coffeebean | Horsenettle | Stinkweed |
| Bedstraw, Pineywoods | Coffeedew | Horseshoe | Stitchwort |
| Beggartick | Coreopsis, Plains | Ironweed | Strawberry, Wild |
| Betony, Florida | Creeping Jenny | Ivy, Poison | Sumac |
| Biden | Croton, Woolly | Jewelweed | Sunflower |
| Bindweed | Cudweed | Jimsonweed | Peppercorn |
| Bindweed, Field | Dandelion | Knapweed, Spotted | Pepperweed, Virginia |
| Bindweed, Hedge | Daisy, English | Knawel | Pigweed |
| Bitter Wintergrass | Daisy, Oxeye | Knawel | Pigweed, Prostrate |
| Bitterness, Hairy | Dandelion, Fall | Knawel | Pigweed, Redroot |
| Bitterweed | Dandelion, False | Knawel | Pigweed, Russian |
| Black-eyed Susan | Dock | Knawel | Pigweed, Smooth Pigweed |
| Bloodflower, Milkweed | Dock, Broadleaf | Knawel | Pigweed, Tumble |
| Brilliant Oxeye | Dock, Curly | Knawel | Pigweed, Tumble |
| Broomweed | Dock, Smooth | Knawel | Pigweed, Tumble |
| Buckhorn | Dogbane | Knawel | Pigweed, Tumble |
| Blackhead, Wild | Dogbane, Hairy | Knawel | Pigweed, Tumble |
| Bullthistle, Wavyleaf | Elderberry | Knawel | Pigweed, Tumble |
| Bur Ragweed | Fiddleneck | Knawel | Pigweed, Tumble |
| Burdock | Fieldress, Austrian | Knawel | Pigweed, Tumble |
| Burweed | Fieldress, Redstem | Knawel | Pigweed, Tumble |
| Buttercup | Flax, False | Knawel | Pigweed, Tumble |
| Buttercup, Bulbous | Flea bane, Daisy | Knawel | Pigweed, Tumble |
| Carpetweed | Flabon, Hairy | Knawel | Pigweed, Tumble |
| Carrot, Wild | Flabon, Rough | Knawel | Pigweed, Tumble |
| Catchweed, Bedstraw | Flaxweed | Knawel | Pigweed, Tumble |
| Camp | Florida Pusley | Knawel | Pigweed, Tumble |
| Catsear | Four-O'Clock, Wild | Knawel | Pigweed, Tumble |
| Catsear, Spotted | Frenchweed | Knawel | Pigweed, Tumble |
| Chamomile, Corn | Galinsoga | Knawel | Pigweed, Tumble |
| Chickweed | Galinsoga, Smallflower | Knawel | Pigweed, Tumble |
| Chickweed, Common | Garlic, Wild | Knawel | Pigweed, Tumble |
| Chickweed, Mouseear | Geranium, Carolina | Knawel | Pigweed, Tumble |
| Chicory | Geranium, Wild Goathead | Knawel | Pigweed, Tumble |
| Cinquefoil | Goldenrod | Knawel | Pigweed, Tumble |
| Cinquefoil, Rough | Ground Ivy | Knawel | Pigweed, Tumble |
| Clematis, Western | Groundsel | Knawel | Pigweed, Tumble |
| Clover | Gumweed | Knawel | Pigweed, Tumble |
| Clover, Annual | Hawweed | Knawel | Pigweed, Tumble |
| Yellow Sweet | Hawweed, Orange | Knawel | Pigweed, Tumble |
| Clover, Crimson | Headal | Knawel | Pigweed, Tumble |
| Clover, Hop | Heartleaf Drymary | Knawel | Pigweed, Tumble |
| Clover, Red | Heathstar | Knawel | Pigweed, Tumble |
| Clover, Strawberry | Hemlock, Poison | Knawel | Pigweed, Tumble |
| Clover, Sweet | | Knawel | Pigweed, Tumble |

USAGE TIPS

- For rectangular lawns**
- Apply two header strips across each end for a turning area, then apply back and forth in the longest direction.
- For irregular-shaped lawns**
- Apply a header strip around the entire lawn for a turning area, then apply back and forth.
- Note:** Each pass should be within 10 feet of each other to ensure an overlap.
- Do not leave spreader open when making turns.
 - For best results, do not apply on a windy day.
 - Swath will vary due to weather conditions, product used and walking speed.
 - Apply products only to lawn.
 - To determine how much coverage you need, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, gardens, walkways, etc.). One bag covers 5,000 sq. ft.



STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. If empty, place in trash or offer for recycling if available. If partly filled, call your local solid waste agency for disposal instructions. Never place unused product down any drain (including toilet) or outdoor (including sewer) drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER OR SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

My Pledge Of Satisfaction

We have made every effort in the formulation of this product to assure that you will be satisfied with the results. If, for any reason, you are not satisfied, please write and describe the problem. Our lawn specialists will make every effort to assure your success.

Distributed & Guaranteed by: Jonathan Green, Inc. POB 326 Farmingdale, NJ 07727

(26002)


(RV032416)

9122

**GREAT
Lawns ONLY
Happen with

PROPER
Soil pH...

and
LOOSE,
Biologically
Active Soil!**



Jonathan Green
QUALITY LAWN SINCE 1881

MAG-I-CAL PLUS®

Natural Soil Food for Lawns in Acidic + Hard Soils

**HELPS
RAISE
SOIL pH**

**HELPS
LOOSEN
HARD SOILS**

**CONTAINS
BIOCHAR**



**20% TO 70% OF YOUR
LAWN FOOD IS WASTED
WHEN SOIL PH IS TOO LOW!**

**PROMOTES AIR AND
WATER PENETRATION
FOR BETTER ROOTING**



FEEDING YOUR SOIL

If you've struggled to grow a great lawn in the past, poor soil is often the problem. If your soil is too acidic (having a pH under 6.2) or compacted, your lawn may not thrive. MAG-I-CAL PLUS for Acidic Soils contains calcium which raises soil pH and improves cell wall structure. Lawns thrive in soil with pH levels between 6.2 and 7, otherwise light green grass and weeds may persist while fertilizer nutrients are not fully used by the grass. MAG-I-CAL PLUS helps to break up clay and compacted soils, which improves root penetration and promotes growth by increasing biological activity while helping release trapped nutrients in the soil.

WHY USE SOIL FOODS?

SOIL pH*: If you are not sure of your soil pH, our soil test kit, available from your local dealer can be helpful. Your local extension office or state university also can provide soil testing.

If the results of your soil test reveal your pH is below 5.9, supplement our New American Lawn program with MAG-I-CAL for Acidic Soils.

Apply any time of the year with other products on the same day. It can be applied when seeding or not.

HARD SOIL:** For hard, compact soil, supplement our New American Lawn Program with LOVE YOUR SOIL.

Apply any time of the year with other products on the same day. It can be applied when seeding or not.

DIRECTIONS FOR USE

MAG-I-CAL PLUS can be applied anytime of the year. Apply only as directed.

ESTABLISHED LAWNS: Apply MAG-I-CAL PLUS using the "Spreader Settings" chart below. For best results, aerate or rake the lawn vigorously prior to applying MAG-I-CAL PLUS. Watering is not necessary, but if possible, water in after application for 15 minutes.

NEW LAWNS OR OVERSEEDING LAWNS OR SODDING: When seeding or sodding lawns, MAG-I-CAL PLUS should be applied with Jonathan Green Green-Up Lawn Food for Seeding & Sodding. These products can be applied in any order, on the same day.

Apply MAG-I-CAL PLUS using the "Spreader Settings" chart below. For best results, aerate or rake the lawn vigorously prior to applying MAG-I-CAL PLUS, grass seed or any of the Jonathan Green products mentioned above. Watering is not necessary, but if possible, water in after application for 15 minutes.

SPREADER SETTINGS

The following settings are approximate. An accurate calculation can be made by application to 1,000 sq. ft. and adjusting spreader opening to apply 3.6 lbs. per 1,000 sq. ft. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .90 pounds per 250 square feet. Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

BROADCAST (ROTARY) SPREADERS

	Established Lawns	New Lawns or pH below 5.9
Jonathan Green		
New American Lawn Standard	4-1/2	Apply
New American Lawn Deluxe & Pro	14	twice over
New American Lawn Hand Held	3	the area
Agri-Fab	2-1/2	the area
Cyclone	4	at the
EZ Republic (Ortho)	4	at the
Earthway (Ames)	13	same
Precision (Red Devil)	5	setting to
Scotts Rotary	5	the left
Scotts EasyGreen	20	the left

DROP SPREADERS

	Established Lawns	New Lawns or pH below 5.9
Agri-Fab	2	Apply
Central	4	twice over
EZ Republic (Ortho)	4	the area
Earthway (Ames)	10	at the
Precision (Red Devil)	3-1/2	same
Scotts	1-1/2	setting to
		the left

For more settings visit www.jonathangreen.com

USAGE TIPS

For rectangular lawns
Apply two header strips across each end for a turning area. Then apply back and forth in the longest direction.

For irregular-shaped lawns
Apply a header strip around the entire lawn for a turning area, then apply back and forth.

Note: For Rotary Spreaders, each pass should be within 10 feet of each other to ensure an overlap.
• Do not leave spreader open when making turns.
• For best results, do not apply on a windy day.
• Swath will vary due to weather conditions, product used and walking speed.
• Apply products only to lawn.
• To determine how much coverage you need, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, gardens, walkways, etc.).

GUARANTEED ANALYSIS

Calcium (Ca)	28.0%
Sulfur (S)	1.5%
1.5% Combined Sulfur (S)	
Iron (Fe)	0.5%
0.01% Water Soluble Iron (Fe)	
Derived from: Calcium Sulfate Dihydrate, Calcium Carbonate, Iron Humate, and Soy Protein Hydrolysate.	

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:
1% Humic Acid (derived from humus)
1% Polyhydroxycarboxylic Acids (derived from plant extracts)
2% Wood Biochar

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

CAUTION

Store in a cool dry place out of the reach of children and pets. Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn or garden and sweep any product that lands on the driveway, sidewalk or street back onto your lawn or garden. Also, sweep fertilizer off concrete surfaces to avoid staining.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all risk of use and/or handling if such use is contrary to label instructions.

CAS NO.

Calcium Sulfate Dihydrate	10101-41-A
Iron Humate	N/A
Calcium Carbonate	1317-65-3
Humic Acid	1415-93-6
Biochar	N/A

Be aware of local ordinance requirements concerning lawn fertilizer applications. "Black out" dates may limit or prohibit lawn fertilizer applications from late fall to early spring in certain areas.

Iowa Secretary of Agriculture certified 1682 pounds of ECCE per ton.



My Pledge Of Satisfaction
We have made every effort in the formulation of this product to insure that you will be satisfied with the results. If, for any reason, you are not satisfied, please write and describe the problem. Our lawn specialists will make every effort to assure your success.



Jonathan Green

Ask for a FREE copy of our
New American Lawn Guide or go to
www.newamericanlawn.com

Sign up to receive timely lawn tips
www.jonathangreen.com

Distributed & Guaranteed by: Jonathan Green, Inc. POB 326 Farmingdale, NJ 07727

21354 SF818



Jonathan Green
QUALITY LAWN SINCE 1881

WINTER SURVIVAL

Fall Lawn Food

HELPS REPAIR
LAWN FROM
SUMMER DAMAGE

IDEAL FOR
FALL
SEEDING

HELPS PROTECT
LAWN FROM
WINTER STRESS

PROMOTES DEEP ROOTING IN FALL...
AND A QUICKER GREEN-UP IN SPRING!

10-0-20

FALL FEEDING

Fall is a great time to grow grass and feed your lawn. Cooler temperatures and higher amounts of rainfall helps your lawn recover from summer stress. Winter Survival Fall Lawn Food provides nutrients to make your lawn thicker and greener than ever before while encouraging deeper-rooting. Winter Survival Fall Lawn Food can be used on all grass types.

HOW TO APPLY

Apply Winter Survival Fall Lawn Food from late summer through fall. For damaged lawns, apply in early fall and again 6 weeks later for a thicker, greener lawn. Apply when grass is dry and water thoroughly after applying for best results; however, product does not require watering in. You may apply MAG-I-CAL Plus in any order on the same day that you apply Winter Survival Fall Lawn Food. You can re-enter lawn area after product is applied. Apply uniformly with a lawn spreader using the chart listed. This package applies 3 lb. of product per 1,000 square feet.

SEEDING LAWNS

Winter Survival Fall Lawn Food is ideal for fall seeding. Apply Winter Survival Fall Lawn Food after grass seed has been planted. You may apply MAG-I-CAL Plus in any order on the same day that you apply Winter Survival Fall Lawn Food. Water the lawn after application to activate the feeding process and get new grass plant seedlings off to a good start.

CARING FOR YOUR LAWN

Mow grass at 2-1/2 to 3-1/2 inches. Do not remove more than 1/3 of the grass blade at one time, with a sharp mower blade. Mulch clippings or leave them on the lawn if not wet or clumpy. One inch of water from rainfall and/or watering per week is usually adequate unless drought conditions exist.

SPREADER SETTINGS

The following settings are approximate for applying contents of this package at a suggested rate of 3 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .75 pounds per 250 square feet. Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

BROADCAST (ROTARY) SPREADERS

Jonathan Green Standard	5
Deluxe & Pro Hand Held	10-11
Agri-Fab	6
Cyclone	5
Earthway (Ames)	14
Lesco	12 - E
Ortho	5
Precision (Red Devil)	5
Scotts Rotary Spreaders	6*
Scotts EasyGreen	26


DROP SPREADERS

Earthway (Ames)	14-1/2
Ortho	5
Precision (Red Devil)	3
Scotts Drop**	7

**Includes Accugreen 1000 - 2000 & Classic


*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 3000

USAGE TIPS



For rectangular lawns

Apply two header strips across each end for a turning area. Then apply back and forth in the longest direction.



For irregular-shaped lawns

Apply a header strip around the entire lawn for a turning area, then apply back and forth.

Note: Each pass should be within 10 feet of each other to ensure an overlap.

- Do not leave spreader open when making turns.
- For best results, do not apply on a windy day.
- Swath will vary due to weather conditions, product used and walking speed.
- To determine how much coverage you need, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, gardens, walkways, etc.).

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	10.00%
10% Urea Nitrogen*	
SOLUBLE POTASH (K ₂ O)	20.00%
IRON (Fe)	3.00%
0.03% water soluble iron	

Derived from: Urea, Polymer Coated Urea, Muriate of Potash, Ferrous Sulfate.

*2.5% Slowly available urea nitrogen from polymer coated urea.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

CAUTION

Keep out of reach of children. Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn or garden and sweep any product that lands on the driveway, sidewalk or street back onto your lawn or garden. Sweep fertilizer off concrete surfaces to avoid staining.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling if such use is contrary to label instructions.

Be aware of local ordinance requirements concerning lawn fertilizer applications. "Black out" dates may limit or prohibit lawn fertilizer applications from late fall to early spring in certain areas.



My Pledge Of Satisfaction
We have made every effort in the formulation of this product to assure that you will be satisfied with the results. If, for any reason, you are not satisfied, please write and describe the problem. Our lawn specialists will make every effort to assure your success.



Ask for a free copy of our Lawn Guide.

Sign up to receive timely lawn tips
www.jonathangreen.com



Distributed & Guaranteed by: Jonathan Green, Inc. POB 326 Farmingdale, NJ 07727

(22400) SF621