


Jonathan Green®


VERI-GREEN®

Southern Weed & Feed For St. Augustine, Zoysiagrass, Centipede & Bermudagrass Lawns

KILLS WEEDS
AND FERTILIZES
THE LAWN

LONG-LASTING
FEEDING

CONTAINS
IRON FOR
DEEP GREENING



DOLLARWEED



DANDELION

KILLS DOLLARWEED
& OVER
40
OTHER WEED TYPES!



SPOTTED SPURGE



WHITE CLOVER

30-0-3

ACTIVE INGREDIENT

penoxsulam: 2-(2,2-difluoroethoxy)-N-(4,5,8-dimethoxy[1,2,4]triazolo[1,5-c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide

0.04%

OTHER INGREDIENTS

99.96%

TOTAL 100.00%

EPA REG. NO. 62719-549-50931
EPA EST. NO. 075527-PA-12 EPA EST. NO. 82757-OH-001 (M)
EPA EST. NO. 82757-MA-002 (F) 8378-IN-1 (K) 69759-OR-004 (LOT BEGINS BAN)
Superscript used is first letter of run code on end of container

For postemergence control of annual and perennial broadleaf weeds in established home lawns and other recreational turfgrass areas. In the state of Florida: Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn/garden.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Refer to back of package for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer and Limitation of Remedies at end of the back of this package. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

GUARANTEED ANALYSIS

Total Nitrogen (N)	30%
30.0% Urea Nitrogen*	
Soluble Potash (K ₂ O)	3%
Iron (Fe)	2%
0.05% water soluble iron (Fe)	

Derived from Urea, Polymer Coated Urea, Muriate of Potash, and Iron Sulfate.

*15% Controlled Release Urea Nitrogen from Polymer Coated Urea

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

F002946

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION:

Causes Moderate Eye Irritation

Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, and shoes plus socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may contact 1-800-992-5994 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in cool dry place in original container.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

HOW THIS PRODUCT WORKS

This product provides postemergence control of annual, biennial and perennial broadleaf weeds in established home lawns and other recreational turfgrass areas.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

- Do not apply this product on dichondra, carpetgrass, bahiagrass, or to turfgrass containing desirable clovers.
- Do not apply this product directly to, or otherwise permit this product to come into contact with, flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur.
- Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at the time of application.
- The active ingredient in this product may leave residues in treated grass clippings which could be harmful to some plants and trees. Do not collect treated grass clippings for use as mulch around plants. The applicator must notify in writing the appropriate residences/individuals of this precaution when this product is applied by a lawn service/commercial applicator.
- To minimize the potential for grass injury, allow four weeks between applications.
- Avoid unnecessary disturbances, including watering and mowing, for approximately 24 hours after application.
- Use this product on soil (monoculture) stands of perennial ryegrass or tall fescue at a maximum rate of 1.2 lb per 1000 sq ft. Some yellowing of these turfgrasses may occur within two weeks of application. Any discoloration is temporary and will dissipate under adequate growing conditions.
- Do not apply more than 3.4 lb of this product per 1000 sq ft per application.
- Do not apply more than 5.1 lb of this product per 1000 sq ft per year.
- Florida Applications: Regardless of grass type, do not apply more than 2.3 lbs. of this product per 1,000 sq. ft.. Do not apply this product more than two times per year. Do not apply to north Florida in winter. Apply only to actively growing turf. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

Maryland Applications: It is a violation of law in Maryland to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq. ft. of readily available nitrogen or more than 0.9 lb per 1000 sq. ft. of total nitrogen with at least 20% of which shall consist of slow release nitrogen per application. Additional restrictions apply to total nitrogen applied to turf per season. Other state and local restrictions may apply.

Virginia Applications: No more than 0.7 lb per 1,000 sq. ft. of readily available nitrogen during any given 30 day period. No more than 0.9 lb (cool season grasses) or 1.0 lb (warm season grasses) per 1,000 sq. ft. of total nitrogen with at least 20% of which shall consist of slow release nitrogen during any given 30 day period.

TREATMENT ON TURFGRASS SPECIES NOT LISTED ON THE LABEL FOR THIS PRODUCT

Users who wish to use this product on turfgrass species not listed on this label may determine the suitability for use by treating a small area at a specified rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the target turfgrass species is tolerant. Applications should not be made to turfgrass under stress. The user assumes responsibility for any plant damage or other liability resulting from use of this product on turfgrass species not listed below on this label.

JONATHAN GREEN VERI-GREEN SOUTHERN WEED & FEED IS RECOMMENDED FOR USE ON THE FOLLOWING ESTABLISHED TURFGRASS SPECIES (numbers in parentheses () refer to footnotes below):

Established Cool Season Turfgrass (1)	Fescue, Sheeps	Centipedegrass (4)
Annual Bluegrass	Fescue, Tall (2)	Fescue, Tall
Bentgrass	Ryegrass, Perennial (2)	(growing in warm season areas) (2)
Bluegrass, Kentucky		Kikuyugrass
Fescue, Chewing	Established Warm Season Turfgrass (3)	Seashore Paspalum
Fescue, Creeping Red	Bermudagrass (common or hybrid)	St. Augustinegrass
		Zoysiagrass

- Use no more than 2.3 lb of this product per 1000 sq ft per application.
- Use this product on these species at a maximum rate of 1.2 lb per 1000 sq ft per application and only if some injury can be tolerated. A slight yellowing of turfgrass may occur within two weeks of application. The discoloration is temporary and is generally gone within four weeks of treatment under adequate growing conditions.
- Use no more than 3.4 lb of this product per 1000 sq ft per application.
- Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at the time of application.

WEEDS CONTROLLED OR SUPPRESSED

Generally, application rates at the lower end of the rate range will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands and/or larger weeds), the higher rate within the rate range or repeat applications will be needed. Weeds growing in the absence of competition from other vegetation generally require higher rates or repeat applications to obtain satisfactory control or suppression.

WEEDS CONTROLLED OR SUPPRESSED			RATE
Burdock	Curly Dock	Lespedeza, Common	Speedwell, Common
Burnweed, American	Dandelion, Common	Matchweed	Spotted Catsear
Buttercup, Hair	Dollarweed	Medic, Black	Spurge, Spotted
Chickweed, Common	Dwarf Beggarweed	Parsley-piert	Spurweed
Chickweed, Mouseear	English Lawn Daisy	Plantain, Broadleaf	Virginia Buttonweed ¹
Clover, Hop	False Dandelion	Plantain, Buckhorn	Virginia Pepperweed
Clover, Red	Florida Betony	Purslane, Common	Wild Violet
Clover, White	Goldenrod	Ragweed, Common	Woodsorrel, Creeping
Cocklebur, Common	Hawkweed	Round Leaf Mallow	Woodsorrel, Yellow
Coffeeweed	Henbit	Sheep Sorrel	
Common Vetch	Ivy, Ground	Shepherd's Purse	
Creeping Beggarweed	Lambsquarters	Smartweed	

¹Kyllinga, Annual and Green (suppression only)¹

¹Nutsedge, Purple and Yellow (suppression only)¹

¹Re-treatment may be necessary

HOW TO APPLY

Apply this product using a drop or rotary-type spreader designed to apply granules. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

To optimize product performance, mow the turfgrass two to three days before applying this product. If the turfgrass is under moisture stress, a thorough sprinkler irrigation of at least 1/2 inch may be needed at least one to two days before application. For optimal results, apply this product in the morning when dew is present. If necessary, sprinkler irrigate with less than 1/4 inch of water just prior to application to maximize particle adherence to leaf surfaces. Avoid watering of treated area for 24 hours following application. Avoid making applications if rainfall is expected within 24 hours after treatment.

APPLICATION TECHNIQUES FOR APPLYING JONATHAN GREEN VERI-GREEN SOUTHERN WEED & FEED

- When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.

SPREADER SETTINGS AS A GUIDE FOR CALIBRATION

Note: These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by Jonathan Green, Inc. or the spreader manufacturer.

BROADCAST (ROTARY) SPREADERS

	2.3 lb. per 1,000 sq. ft.
Jonathan Green Standard Rotary Deluxe Hand Held	5-1/2 14 3
Agri-Fab	6-1/2
Cyclone	5
Earthway (Ames)	13
Echo RB60	6-1/2
EZ Republic (Ortho)	3
Lesco	11 - D-1/2
Precision (Red Devil)	5
Scotts Rotary Spreaders*	6-1/2
Spyker	5 - 6
STA-GREEN	5
Vigoro	6-1/2

*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 3000

DROP SPREADERS

	2.3 lb. per 1,000 sq. ft.
Agri-Fab	7
Central	7
EZ Republic (Ortho)	7
Earthway (Ames)	13
Precision (Red Devil)	1-1/2
Scotts Drop **	4 - 5

**Includes Accugreen 1000 - 2000 & Classic

For more settings visit www.jonathangreen.com

HOW MUCH TO APPLY

Apply this product to established turfgrass. Possible re-treatment in approximately 30 days may be necessary if turfgrass is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed specified rates because damage to turfgrass may occur. Do not make more than two applications per season.

Note: When applying to warm season turfgrasses under conditions of inadequate moisture, high temperature and high humidity, over application may cause minor temporary discoloration. It is recommended to reduce application rates by 50% during hot, dry conditions to avoid possible dehydration or browning of turfgrass.

WHEN TO APPLY

Best weed control results are obtained when this product is applied to small, actively growing weeds in the spring or early fall. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control. Broadleaf weed species germinate at different times.

This product may also be applied to newly established warm season turfgrass. Delay treatments four weeks after sodding, sprigging or plugging.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- Refund of purchase price paid by buyer or user for product bought, or
- Replacement of amount of product used.

To the extent permitted by law, Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

To the extent permitted by law, the terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.